

TimTec Diversity Library of 10,000 compounds was screened in research collaboration between Department of Veterinary Molecular Biology at Montana State University, Bozeman, Montana, and Department of Chemistry at Altai State Technical University, Barnaul, Russia.

A screen of the library identified 26 test compounds and explored their effects on human neutrophil function.

Schepetkin, I., et al., High-throughput Screening for Small-molecule Activators of Neutrophils: Identification of Novel N-Formyl Peptide Receptor Agonists. *Mol. Pharmacol.* 71: 1061-1074 (2007)

Abstract

We screened a chemolibrary of drug-like molecules for their ability to activate reactive oxygen species (ROS) production in murine phagocytes, and we identified 26 novel compounds with potent neutrophil activating properties. We used substructure screening, fragment-focusing, and structure-activity relationship analyses to further probe the parent library and defined at least two groups of activators of ROS production in murine neutrophils: t-butyl benzene and thiophene-2-amide-3-carboxylic ester derivatives. Further studies of the active compounds revealed 11 compounds that activated ROS production in human neutrophils, and six of these compounds also activated intercellular Ca²⁺ mobilization and chemotaxis in human neutrophils. Of the latter compounds, compound 14 (1,3-benzodioxolane-5-carboxylic acid 4[1]-benzyloxy-3[1]-ethoxybenzylidene-hydrazide) activated neutrophils at nanomolar concentrations, and Ca²⁺ mobilization was inhibited by pertussis toxin and N-t-butoxycarbonyl-Phe-Leu-Phe-Leu-Phe (Boc-2), an antagonist of formyl peptide receptors (FPR/FPRL1). Likewise, activation by compound 14 was desensitized after N-formyl-Met-Leu-Phe pretreatment. Similar biological activities were found for compound 104 (1,3-benzodioxolane-5-carboxylic acid 3[1]-bromo-5[1]-ethoxy-4[1]-hydroxybenzylidenehydrazide), an analog of compound 14. Furthermore, conformational analysis of the activators of chemotaxis and Ca²⁺ mobilization showed a high degree of similarity in distances between pharmacophore points of compounds 14 and 104 with a model of FPR published by Edwards et al. (*Mol Pharmacol* 68:1301–1310, 2005), indicating that conformational features of the agonists identified here are structurally compatible with steric constraints of the ligand-binding pocket of the receptor. Based on these results, we conclude that compounds 14 and 104 represent novel small-molecule agonists of FPR. These studies enhance our understanding of FPR ligand/receptor interactions and structure/activity relationships of phagocyte agonists.

[Activators of Neutrophils analogs](#)

Activators of Neutrophils

ST044918

C₁₇H₂₀N₄O₅S

392.44

C₁₅H₁₈N₂O₅


306.32



C₂₂H₂₃CIN₆O

440.54

ST032745


ST031779







~~ST00588A~~


ST00588A







~~ST028770~~


ST028770







~~ST000299C~~


ST000299A







~~XXXXXXXXXX~~


~~XXXXXXXXXX~~








~~ST020970~~

ST020970





BRACMA283AM5QS



[Activators of Neutrophils analogs](#)